



Identification of Bioactive Components in the Hydroalcoholic Extract of *Syringodium Isoetifolium* and Assessment of its Biological Activity by Gas Chromatography – Massspectrometry

Kalaivani P, Amudha P*

Department of Biochemistry, Vels Institute of Science, Technology and Advanced Studies, Pallavaram, Chennai-600117, Tamil Nadu, India



Article History:

Received on: 22 Jan 2021
Revised on: 26 Feb 2021
Accepted on: 17 Mar 2021

Keywords:

Syringodium
isoetifolium,
Hydroalcoholic extract,
Gas Chromatography and
Mass Spectrometry
Analysis,
Octadecanoic acid,
Biological activities

ABSTRACT

Herbal plants show a growing interest in natural antioxidant, also play an important role in the treatment of muscle pain, stomach problems, wounds etc. About 75 to 80% of the world population uses herbal medicine. One such herb (Marine Plants) used in the treatment of cancer is *Syringodium isoetifolium*. In a previous study, *Syringodium isoetifolium* with three different extracts (Aqueous, ethanol and 70% hydroalcohol) proves the presence of phytochemical such as Tannin, Saponin, Flavonoids, Steroids, Terpenoids, coumarin etc and also has high antioxidant activity. The current study aims to identify the bioactive compounds from the hydroalcoholic extract of *Syringodium isoetifolium*. The photo components present in the hydroalcoholic extract was determined by GC-MS Analysis and the mass spectra of compounds present in the extract were matched with the National Institute of Standards and Technology library. 20 different phytochemicals were found to be present in the hydroalcoholic extract of *Syringodium isoetifolium* using GC-MS analysis. Some of the important phytochemicals are Hexanoic acid, Methyl ester, Octadecanoic acid, 1,2-Benzenedicarboxylic acid, bis(2-methylpropyl) ester, 1-Isopropylidene-3-methyl-3-vinylcyclobutane. The study reveals that the identified phytochemicals from the hydroalcoholic extract of *Syringodium isoetifolium* has essential biological activities such as Antibacterial, Antifungal, Antimicrobial, Antifouling and Anticancer properties. Further depth of the study will deal with the exact mode of action of the phytochemicals.

*Corresponding Author

Name: Amudha P
Phone: 7904075676
Email: amudhaa85@gmail.com

ISSN: 0975-7538

DOI: <https://doi.org/10.26452/ijrps.v12i2.4650>

Production and Hosted by

IJRPS | www.ijrps.com

© 2021 | All rights reserved.

INTRODUCTION

From the ancient period, natural herbs play a significant role in the treatment of diseases such as fever, stomach problems, muscle pains, wounds. But nowadays, we forget about traditional foods, as well as natural medicine. The marine sources are potent therapeutic drugs for various ailments (Lee and Jeon, 2013). Under the stress condition, the compounds present in the seagrass produce a defence mechanism due to the production of secondary metabolites (Subhashini et al., 2013). One such seagrass, *Syringodium isoetifolium*, commonly called as Noodle grass, Neer pasi, Oosi korai, Nool pasi, Korai pasi, were found in the subtidal region, especially




Figure 1: GC-MS Chromatogram of the Hydro-alcoholic extract of *Syringodium isoetifolium* by GC-MS Analysis

in India richly found in the Andaman and Lakshadweep islands (Kalaivani *et al.*, 2020).

Secondary metabolites such as flavonoids, alkaloids, terpenoids, tannins and steroids were present in *Syringodium isoetifolium*, which can be supplemented as a traditional medicine to human beings. The phytochemicals present in the seagrass *Syringodium isoetifolium* can be separated and identified by GC-MS Analysis. The phytochemicals can be interpreted and matched the spectrum with the National Institute of Standard and Technology (NIST) (Amudha *et al.*, 2018). The best technique to identify the long chain hydrocarbons, esters alkaloids, alcohols, nitro and amino compounds (Jadhav *et al.*, 2014).

The present study deals with the phytochemicals present in the hydroalcoholic extract of *Syringodium isoetifolium* by GC-MS analysis and biological activities of the compounds were determined.

MATERIALS AND METHODS

Collection of seagrass

The fresh seagrass *Syringodium isoetifolium* were collected from Devapattinam, Ramanathapuram District, on June 2019 by sea divers. The sample has been identified and authenticated by Dr P.Jeyaraman, PhD., Director, Retd Professor, Presidency College, in Plant Anatomy Research Centre. The collected seagrass was washed with the tap water, shadow dried and powdered in the herbal grinder. The powdered sample was stored in an airtight container.

Extraction

With three different conical flasks, 20 grams of powdered sample were added and the flask contains different solvent such as aqueous, ethanol and 70% hydroalcohol in the measurement of 1000ml. By the rotator shaker, they were shaken vigorously for one hour and the mixture was kept for 24

Table 1: Identification of Bioactive components in the Hydro-alcoholic extract of *S.isoetifolium* using GC-MS Analysis

Peak	Retention Time	Area %	Height %	Molecular Formula	Molecular Weight	Name of the compounds
1	5.917	4.91	5.12	C ₁₀ H ₁₆	136	1-Isopropylidene-3-methyl-3-vinylcyclobutane
2	6.958	2.00	2.03	C ₃ H ₄	40	1-propyne (cas) propyne
3	8.143	1.85	1.46	C ₃ H ₄	40	1,2-propadiene (cas) allene
4	8.358	2.14	1.55	C ₃ H ₄	40	1-propyne (cas) propyne
5	10.375	1.26	1.48	C ₁₇ H ₂₆ O ₂	262	1,1'-bibicyclo(2.2.2)octyl-4-carboxylic acid
6	13.133	1.64	2.75	C ₂₀ H ₂₆ O ₄	330	1,2-Benzenedicarboxylic acid, dicyclohexyl ester
7	13.333	1.48	1.79	C ₃ H ₄	40	1-propyne (cas) propyne
8	13.578	2.09	3.72	C ₇ H ₁₄ O ₂	130	Hexanoic acid, methyl ester
9	13.691	4.18	6.75	C ₁₄ H ₂₂ N ₂ O	234	Xylocaine
10	13.998	29.99	31.73	C ₁₈ H ₃₆ O ₂	284	Octadecanoic acid
11	14.150	3.46	4.20	C ₁₆ H ₂₂ O ₄	278	1,2-Benzenedicarboxylic acid, bis(2-methylpropyl) ester
12	14.220	2.29	3.07	C ₈ H ₁₁ NO	137	7-hydroxy-5,6,7,8-tetrahydroindolizaine
13	14.345	3.03	4.20	C ₁₅ H ₃₀ O ₂	242	Oxirane, [(dodecyloxy)methyl]
14	15.417	1.59	1.69	C ₂₂ H ₁₃ NO ₄	355	Ethyl 1-hexyl-4-hydroxy-2(1h)-oxo-3-quinolinecarboxylate
15	15.575	3.05	1.93	C ₂₂ H ₁₃ NO ₄	355	Ethyl 1-hexyl-4-hydroxy-2(1h)-oxo-3-quinolinecarboxylate
16	15.883	24.50	17.53	C ₁₈ H ₃₄ O ₂	282	9-Octadecenoic acid
17	16.108	3.18	2.16	C ₁₂ H ₁₂ F ₆ O ₂	302	8,9,9,10,10,11-hexafluoro-4,4-dimethyl-3,5-dioxatetracyclo[5.4.1.0(2,
18	16.493	2.74	3.48	C ₁₅ H ₃₀ O ₂	242	Oxirane, [(dodecyloxy)methyl]-
19	22.606	2.52	2.21	C ₁₅ H ₁₄ N ₂ O ₂	254	4-(methoxymethyl)-6-methyl-2-phenoxy nicotinonitrile
20	24.235	2.08	1.14	C ₃ H ₄	40	1-propyne (cas) propyne

S.isoetifolium: *Syringodium isoetifolium*, GC-MS: Gas chromatography-Mass spectrometry

hours. Then the different extracts were filtered through Whatman no.1 filter paper and the filtrate used for further studies. Hydroalcoholic extract of *Syringodium isoetifolium* shows more Secondary Phytochemicals in the previous preliminary studies and hence it has been used in this study.

GC-MS Analysis

GC-MS analysis was carried out following conditions: column RTX 5Ms with column diameter is 0.32mm; column thickness 0.50 μ m, column length is 30m, operating in electron impact mode at 70eV. Carrier gas used in this analysis is the helium gas (99.999%) with a constant flow of 1.73 ml/min. Injection volume employed was 0.5 μ l was employed (split ratio of 10:1) with an injector tem-

perature of 270°C and an ion-source temperature is 200°C. The temperature of the column was programmed from 40°C (isothermal for 2 min), with an increase of 8°C/min, to 150°C, then 8°C/min to 250°C and finally ends with 20min isothermal at 280°C. At 70eV, the Mass spectra were measured at an interval of 0.5 seconds and fragments from 40 to 450 Da. The total running time of GC are 51.25min. The relative percentage amount of each component was calculated by comparing its average peak area to the total areas. The phytochemicals were identified with the software adopted to handle mass spectra. And chromatograms was a Turbo Mass Ver 5.2.0 (Srinivasan *et al.*, 2013).

Identification of components

Table 2: Biological activities of the identified compounds in Hydro-alcoholic extracts of *S.isoetifolium* by GC-MS Analysis

Peak	R. Time	Name of the compounds	Biological activities
1	13.998	Octadecanoic acid	Cosmetic, Flavor, Hypocholesterolemic, Lubricant, Perfumery, Propepic, Suppository, Lower LDL Cholesterol level Kanthal et al. (2014)
2	15.883	9-Octadecenoic acid	Antihypertensive, Allergenic, Anti-inflammatory, Anticancer, Antiandrogenic, Flavour, Irritant Increase HDL and decrease LDL Cholesterol, Velayutham and Karthi (2015)
3	13.578	Hexanoic acid, methyl ester	Antibacterial, Antifungal
4	14.150	1,2-Benzenedicarboxylic acid, bis(2-methylpropyl) ester	Antimicrobial and Antifouling Kanthal et al. (2014)
5	22.606	4-(methoxymethyl)-6-methyl-2-phenoxynicotinonitrile	Anticancer, Mucolytics, Drugs for the genital disorder Tyagi and Agarwal (2017)
6	15.417	Ethyl 1-hexyl-4-hydroxy-2(1h)-oxo-3-quinolinecarboxylate	Antimicrobial
7	16.493	Oxirane	Natural mediators for allergic Asthma Rao et al. (1983)
8	13.333	1-propyne (cas) propyne	Antidepressive or Antihypertensive agents Maycock et al. (1976)

S.isoetifolium:*Syringodium isoetifolium*,GC-MS: Gas chromatography-Mass spectrometry

The phytochemicals present in the hydro alcoholic extract of *Syringodium isoetifolium* was identified by comparing spectrum with a database of National Institute Standard and Technology (NIST) having more than 62,000 patterns. The name of the compound, molecular formula, structure and also the retention time were determined. The spectrum of the unknown and known components was compared and stored in the NIST library.


RESULTS AND DISCUSSION

Identification of Bioactive components in Hydroalcoholic extract of *Syringodium isoetifolium* by GC MS Analysis


The compounds such as esters, alkaloids, steroids and alcohols present in the test sample can be identified by the combination of Gas Chromatography and Mass Spectrometry technique ([Saravanan et al., 2014](#)). GC-MS also used to analyse a specific test that can identify the forensic substances ([Wagner et al., 1984](#)). Twenty compounds were identified in the hydro-alcoholic extract of *Syringodium isoetifolium* by GC-MS analysis. The active principles with their concentration (%), retention time (RT), molecular

formula and molecular weight (MW) are presented in Table 1. The chromatogram of the hydro-alcoholic extract of seagrass *Syringodium isoetifolium* was shown in Figure 1.


The prevailing compounds are Octadecanoic acid, Hexanoic acid methyl ester, 1,2-Benzenedicarboxylic acid, bis(2-methylpropyl) ester and 9-Octadecenoic acid were found in seagrass *Syringodium isoetifolium*. Octadecanoic acid shows the highest peak with a retention value of 13.998 when compared to the other compounds. And the second peak value was shown by the compound 9-Octadecenoic acid (Oleic acid) with the retention time of 15.883. The other compounds are 1-Isopropylidene-3-methyl-3-vinylcyclobutane (5.917), 1-propyne (cas) propyne (6.958), 1,2-propadiene (cas) allene (8.143), Xycaine (13.691), Oxirane, [(dodecyloxy)methyl (14.345). However, isolation of the individual phytochemical constituents and subjecting to its biological activity will give fruitful results. The biological activities of the compounds present in the hydro-alcoholic extract *Syringodium isoetifolium* was given in Table 2 and the structure of some of the compounds was shown in Figure 2.





Octadecanoic acid





9-Octadecenoic acid



Hexanoic acid, methyl ester

1,2-Benzenedicarboxylic acid,
bis(2-methylpropyl) ester4-(methoxymethyl)-6-methyl-2-
phenoxynicotinonitrile

1-propyne (cas) propyne

**Figure 2: Structure of the compounds identified by GC-MS analysis**

Octadecane are the long chain alkanes and acts as a lubricant and anticorrosion agents. The highest peak value shown by Octadecanoic acid has the following biological activities as Antimicrobial, Anticancer, Hepatoprotective, Anti-arthritis, Anti-asthma and Diuretic (Kanthal *et al.*, 2014). 4-(methoxymethyl)-6-methyl-2-phenoxynicotinonitrile has anticancer activity and they also used as a drug for the genital disorder. 2-Benzenedicarboxylic acid, diethyl ester was used in Plasticizers. 1,2-Benzenedicarboxylic acid reduces oxidative stress and able to cure neurodegenerative disorders (Tyagi and Agarwal, 2017). Oxirane are the natural mediators for allergic asthma and they also act as a biogenetic precursor of leukotrienes (Rao *et al.*, 1983). Oxirane also exhibits antibacterial, insect antifeedant

and antioxidant activities (Thirunarayanan and Vanangamudi, 2011). The bioactive compound 1-propyne (cas) propyne acts as antidepressive or antihypertensive agents (Maycock *et al.*, 1976). Xylocaine compound has anti-pyretic activity and also acts as a local anesthetic (Chakraborty *et al.*, 2010). 9-Octadecenoic acid has many biological activities such as Antiandrogenic, Allergenic, Anti-inflammatory, Antileukotrienes, Anticancer (Velayutham and Karthi, 2015). This paper reveals the importance and goodness of *Syringodium isoetifolium*. Based on the phytochemicals present, the biological activities of earlier studies have been reported. Hence the *Syringodium isoetifolium* is a natural medicine without any side effects and also cost effective.

CONCLUSIONS

In the present investigation, nearly twenty compounds have been identified from the hydroalcoholic extract of *Syringodium isoetifolium* using GC-MS analysis. The presence of bioactive compounds in the seagrass *Syringodium isoetifolium* proved pharmaceutical importance. The various bioactive compounds present in the hydroalcoholic extract of *Syringodium isoetifolium* justify its use for various ailments by traditional practitioners. Further studies and investigation may proceed with the mode of action of each compound.

ACKNOWLEDGEMENT

I am thankful to the Department of Biochemistry, School of Lifesciences (VISTAS) for providing me an opportunity to publish my research paper.

Funding Support

The authors declare that they have no funding support for this study.

Conflict of Interest

The authors declare that they have no conflict of interest for this study.

REFERENCES

- Amudha, P., Jayalakshmi, M., Pushpabharathi, N., Vanitha, V. 2018. Identification of Bioactive Components in *Enhalus acoroides* seagrass extract by Gas Chromatography- Mass Spectroscopy. *Asian Journal of Pharmaceutical and Clinical Research*, 11(10):313-317. ISSN: 0974-2441, 2455-3891.
- Chakraborty, A., Devi, B. R. K., Sanjebam, R., Khumbong, S., Thokchom, J. S. 2010. Preliminary studies on local anesthetic and antipyretic activities of *Spilanthes acmella* Murr. in experimental animal models. *Indian Journal of Pharmacology*, 42(5):277-277. ISSN: 0253-7613.
- Jadhav, V., Kalase, V., Patil, P. 2014. GC-MS analysis of bioactive compounds in methanolic extract of *Holigarna grahamii* (wight) Kurz. *International Journal of Herbal Medicine*, 2(4):31-35.
- Kalaivani, P., Kavitha, D., Vanitha, V. 2020. A Review on Phytochemical and Pharmacological activities of *Syringodium isoetifolium*. *International Journal of Research in Pharmaceutical Sciences*, 11(1):207-214. ISSN: 0975-7538.
- Kanthal, L., Dey, A., Satyavathi, K., Bhojaraju, P. 2014. GC-MS analysis of bio-active compounds in methanolic extract of *Lactuca runcinata* DC. *Pharmacognosy Research*, 6(1):58-61. ISSN: 0974-8490.
- Lee, S.-H., Jeon, Y.-J. 2013. Anti-diabetic effects of brown algae derived phlorotannins, marine polyphenols through diverse mechanisms. *Fitoterapia*, 86:129-136. ISSN: 0367-326X.
- Maycock, A. L., Abeles, R. H., Salach, J. I., Singer, T. P. 1976. The structure of the covalent adduct formed by the interaction of 3-dimethylamino-1-propyne and the flavine of mitochondrial amine oxidase. *Biochemistry*, 15(1):114-125. ISSN: 0006-2960, 1520-4995.
- Rao, A. S., Paknikar, S. K., Kirtane, J. G. 1983. Recent advances in the preparation and synthetic applications of oxiranes. *Tetrahedron*, 39(14):2323-2367. ISSN: 0040-4020.
- Saravanan, P., Chandramohan, G., Mariajancyrani, J., Shanmugasundaram, P. 2014. GC-MS analysis of Phytochemical Constituents in Ethanolic bark extract of *Ficus religiosa* Linn. *International Journal of Pharmacy and Pharmaceutical Sciences*, 6(1):457-460.
- Srinivasan, K., Sivasubramanian, S., Kumaravel, S. 2013. Phytochemical profiling and GC-MS study of *Adhatoda vasica* leaves. *Int.J.Pharm.Bio.Sci*, 5(1).
- Subhashini, P., Dilipan, E., Thangaradjou, T., Papenbrock, J. 2013. Bioactive natural products from marine angiosperms: abundance and functions. *Natural Products and Bioprospecting*, 3(4):129-136. ISSN: 2192-2195, 2192-2209.
- Thirunarayanan, G., Vanangamudi, G. 2011. Synthesis, spectral studies, antimicrobial, antioxidant and insect antifeedant activities of some 9H-fluorene-2-yl keto-oxiranes. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 81(1):390-396. ISSN: 1386-1425.
- Tyagi, T., Agarwal, M. 2017. Phytochemical screening and GC-MS analysis of bioactive constituents in the ethanolic extract of *Pistia stratiotes* L, and *Eichhornia crassipes*. *Journal of Pharmacognosy and Phytochemistry*, 6(1):195-206.
- Velayutham, P., Karthi, C. 2015. Gc-Ms Profile of invitro, invivo and fungal elicited invitro leaves of *Hybanthus enneaspermus* (L.)F.Muell. *International Journal of Pharmacy and Pharmaceutical Sciences*, 7(10):260-267.
- Wagner, H., Bladt, S., Zgainski, E. M. 1984. Plant Drug Analysis. pages 298-334, Berlin. Springer- Verlag. ISBN: 978-3-642-00574-9.