ORIGINAL ARTICLE



INTERNATIONAL JOURNAL OF RESEARCH IN PHARMACEUTICAL SCIENCES

Published by JK Welfare & Pharmascope Foundation

Journal Home Page: <u>www.ijrps.com</u>

Phytochemical Screening and *In Vitro* Antibacterial Activities of Leaf Extract *Acacia etbaica Schweinf* Against Multidrug Resistant Enterobacteriaceae Human Pathogens

Tekleweyni Tadege¹, Kidu Hintsa¹, Tesfay Weletnsae¹, Gopalakrishnan V K¹, Muthulingam M², Kamalakararao K³, Krishna Chaithanya K^{*1}

¹Department of Chemistry, College of Natural and Computational Sciences, Aksum University, Axum, Tigray Region, Ethiopia

²Department of Zoology, Faculty of Science, Annamalai University, Chidambaram, Tamil Nadu, India ³Department of Biochemistry, University College of Science and Technology, Adikavi Nannaya University, Rajamahendravaram, Andhra Pradesh, India

Article History:	ABSTRACT
Received on: 22 Apr 2020 Revised on: 13 May 2020 Accepted on: 15 Jun 2020 <i>Keywords:</i>	Acacia etbaica Schweinf belongs to the family Fabaceae widely distributed in Africa and various parts of this plant such as bark, leaves, flowers and roots are widely used as a folk medicine for curing of various ailments. This study was aimed to screen the phytoconstituents and evaluate the antibacterial activ-
Acacia etbaica Schweinf, Antibacterial agents, Enterobacteriaceae family, Agar well diffusion, Minimum inhibitory concentration	Ity of fear extract of <i>A. etbaica</i> against selected multiding resistant <i>Enterobacteriaceae</i> family. Leaves of <i>A. etbaica</i> were extracted with petroleum ether, chloroform, acetone, and ethanol by sequential soxhlet extraction. <i>In vitro</i> antibacterial activities of <i>A. etbaica</i> leaf extracts against selected <i>Enterobacteriaceae</i> family gram-positive bacteria such as (<i>B. subtilis, E. faecalis, S. aureus</i>) and gram-negative (<i>E. coli, K. pneumonia, V. cholera</i>) were evaluated by agar well diffusion. The antibacterial potential of acetone and ethanol leaf extracts of <i>A. etbaica</i> were determined by 96 well plate broth dilution assay. Among the tested organic leaf extracts, both acetone and ethanolic leaf extract of <i>A. etbaica</i> showed a potentially broad spectrum of <i>in vitro</i> antibacterial activity against tested multiple drug resistant <i>Enterobacteriaceae</i> family grampositive pathogens such as <i>B. subtilis, E. faecalis</i> , and <i>S. aureus</i> and gramnegative bacterial strains <i>E. coli, K. Pneumonia</i> and <i>V. Cholera</i> with significant MIC values. The significant antibacterial activity of both acetone and ethanolic leaf extracts of <i>A. etbaica</i> was due to the subsistence of secondary metabolites phytoconstituents such as flavonoids and phenolic compounds.

*Corresponding Author

Name: Krishna Chaithanya K Phone: +251 944121156 Email: krishanchaitanyawc@gmail.com

ISSN: 0975-7538

DOI: <u>https://doi.org/10.26452/ijrps.v11i3.2782</u>

Production and Hosted by

IJRPS | www.ijrps.com

@ 2020 \mid All rights reserved.

INTRODUCTION

According to CDC multiple and extensive drug resistant (antibiotic-resistant) clinical bacterial strains causes approximately 2.6 million antibioticresistant infections and about 44,000 deaths occurred each year, and the mortality rate of antibiotic resistance is gradually it becomes two-fold each year (CDC-Antibiotic Resistance Threats in the United States, 2019). (Reta *et al.*, 2019) from his cross-sectional study on antibiotic-resistant at tertiary care hospitals at Hawassa University Comprehensive Specialized Hospital in Southern

Phytochemicals	Acacia etbaica							
	Petroleum ether	Chloroform	Acetone	Ethanol				
Alkaloids	-	-	-	-				
Flavonoids	-	+	+++	++				
Coumarins	+	+	+	-				
Tannins	_	+	+++	++				
Terpenoids	+	+	+++	++				
Phenolic compounds	-	+	+++	++				

Table 1: Phytochemical screening of Acacia etbaica

NB: (+) Slightintensity; (++)Moderate intensity; (+++)High intensity; (-)Absent

Table 2: Zone of inhibition (mm) of petroleum ether leaf extract Acacia etbaica (PEAE)

Extract/ Standard	Concen tration (µg/mL)	Zone of inhibition (mm)					
		B.substilis	S.aureus	E. faecalis	K.pneumon	V. cholera	E.coli
AEPE	100	NA	NA	NA	NA	9	10
	250	NA	NA	NA	NA	11	11
	500	NA	NA	10	NA	13	13
Ciproflaxin	20	36	33	36	33	31	26

Table 3: Zone of inhibition (mm) of chloroform extract Acacia etbaica (CHAE)

Extract/	Concen		Zone of inhibition (mm)					
standard	tration				· · ·			
Stanuaru	tration							
	$(\mu g/mL)$							
		B.substilis	S.aureus	E. faecalis	K.pneumonia	V. cholera	E.coli	
AECH	100	NA	10	10	NA	NA	9	
	250	11	11	11	11	11	12	
	250	11	11	11	11	11	12	
	500	12	12	12	12	12	13	
Cinroflavin	20	36	33	36	33	31	26	
ыргоналш	20	50	55	50	55	51	20	

Table 4: Zone of inhibition (mm) of acetone leaf extract Acacia etbaica (ACAE)

		()				()		
Extract/	Concen	Zone of inhibition (mm)						
standard	tration							
	(μ g/mL)							
		B.substilis	S.aureus	E. faecalis	K.pneumon	V. cholerae	E.coli	
AEAC	100	12	12	11	11	10	10	
	250	14	13	13	13	12	15	
	500	16	15	15	15	16	17	
Ciproflaxin	20	36	33	36	33	31	26	

Extract/ standard	Concen tration (µg/mL)	Zone of inhibition (mm)					
		B.substilis	S.aureus	E. faecalis	K.pneumon	V.cholerae	E.coli
AEET Ciproflaxin	100 250 500 20	10 12 14 36	10 11 14 33	10 12 14 36	10 12 14 33	11 12 13 31	NA 11 18 26

Table 5: Zone of inhibition (mm) of ethanol leaf extract Acacia etbaica (ACET)

Table 6: Minimum Inhibitory Concentration (MIC) of acetone and ethanolic leaf extract of *Acacia etbaica* on Enterobacteriaceae family

Plant	MIC (μ g/ml) of acetone and ethanolic leaf extract of <i>Acacia ethaica</i> against three gram								
Extract	positive and three gram negative bacteria								
	B.subtilis	E.feacalis	S.aureus	E.coli	K.Pneumonia	V. cholera			
AE.AC	0.53	2.58	0.63	1.56	4.12	2.06			
AE.ET	1.24	4.12	1.5	3.01	24.87	14.86			

Ethiopia, reported that among the gram-positive bacteria isolates *S.aureus* and gram-positive bacteria *Klebsiella spp* are more prevalence, the gram +ve bacterial isolates showed 81.8 % and 81.1 % antibiotic resistance against penicillin G, cotrimoxazole respectively, and gram -ve bacterial isolates showed antibiotic resistance to 85 % and 93.1% to tetracycline and cotrimoxazole respectively.

Diarrheal diseases caused by E.coli and V.cholera have been an important public health issue of developing countries like Ethiopia leading to high morbidity and 13.6% mortality under-five children (Melese et al., 2019). Enterobacteriaceae family such as several strains of E.coli, Klebsiella, Enterobacter and Salmonella spp. showed high drug-resistant against ampicillin, cotrimoxazole and tetracycline respectively and responsible for pathogenesis of nosocomial and community-acquired infections such as urinary, enteric, wound, burn, bloodstream and pneumonia infections (Brenner et al., 2005). For the last two decades, some of the clinically important pathogenic bacterial strains including the Enterobacteriaceae family develop multiple and extensive antibiotic resistance (MDR and XDR) against standard Synthetic, semi-synthetic antibiotics and spreading of resistance among the clinically bacterial strains and gradually standard antibiotics losing their efficacy for treating communicable diseases (Fair and Tor, 2014).

The bacterial strains showed induced antibacterial resistance to antibiotics in the pinnacle of evolution (Munita and Arias, 2016). Due to the genetic plasticity of bacterial pathogens that trigger neutralization of antibiotic effect, transition of binding receptor site of antibiotics and attainment of genes coded for MDR and XDR (Nikaido, 2009). The emerging of bacterial strains to withstand the effect of antibiotics and undesirable secondary effects associated with synthetic antibacterial agents, microbiologists search for potential antimicrobial substances from various sources of natural sources for combating infectious diseases. In Ethiopia, since long back traditional practicing of medicinal plants has become an integral part of the culture that has been used to treat different human infectious diseases (Jima and Megersa, 2018).

Acacia etbaica Schweinf belongs to the family *Fabaceae* widely distributed in Ethiopia, Eritrea, Kenya, Somalia, Sudan, Tanzania, and Uganda (Ross, 1979).

Various plant parts of *A.etbaica* such as bark, flowers, leaves, pods, seeds, and roots are commonly used by traditional healers to cure various ailments (Kassem *et al.*, 2014). The leaves of *A.etbaica* are applied as a folk remedy to heal swelling by crushing the leaves, mixed with the latex of *Erythrina abyssinica* and rub the paste on the affected part of the swelling area and also used at infected part the body (Teklay *et al.*, 2013).

Hence the current study was intended to evaluate the *in vitro* antibacterial activity organic leaf extracts of *A. etbaica* against the selected multiple



Figure 1: Zone of inhibition (mm)of petroleum ether leaf extract of *Acacia etbaica* (100, 250, 500μ g/mL) and standard antibiotics Ciproflaxin (20 μ g/mL) against three gram negative and three gram positive bacteria by the Agar well-diffusion method

drug-resistant Enterobacteriaceae family.

MATERIALS AND METHODS

Description of study areas

The plant sample was collected from Axum city in the Central Zone of Tigray Ethiopia. The phytochemical analysis and antibacterial potential of leaf extract of *Acacia ethaica* was carried out in the Chemistry Laboratory, Aksum University, Tigray region, Ethiopia.

Sample collection and authentication

Leaves of *A.etbaica* were collected from Axum city in January 2018 and transported to the Chemistry Laboratory at Aksum University for further processing. The plant specimen was authenticated by Mr. Melaku Wendafrash (M.Sc.) National Herbarium, Department of Biology, Addis Ababa University, Addis Ababa, and it was given the voucher spec-

imen number TT 001.

Preparation of Plant extract

The leaves of *A.etbaica* were properly washed with distilled water, dried in shade at room temperature for two weeks days, crushed into fine powder by using an electrical grinder. 100 grams of plant powder was taken and subjected to sequential extraction by Soxhlet apparatus using 500ml of petroleum ether, chloroform, acetone & ethanol, and these organic extracts were concentrated at 35-40°C under reduced pressure and vacuum using a rotary evaporator (Das *et al.*, 2010). The concentrated plant extracts were then put in an airtight container and stored at 4° C before use in phytochemical and antibacterial activities.

Phytochemical screening of organic leaf extract

The phytochemical screening of *A. etbaica* leaf organic extracts were conducted to determine the presence of phytoconstituents like alkaloids,



Figure 2: Zone of inhibition (mm) of chloroform leaf extract of *Acacia etbaica* (100, 250,500 μ g/mL) and standard antibiotics Ciproflaxin (20 μ g/mL) against three gram negative and three gram positive bacteria by the Agar well-diffusion method

flavonoids, tannins, coumarins, terpenoids, and phenols (Harborne, 1998).

Bacterial Strains

The bacterial strains such as three Gram-positive *Bacillus subtilis* (ATCC3915) *Enterococcus faecalis* (ATCC29212), *Staphylococcus aureus* (ATCC25923) and three Gram-negative bacteria *Escherichia coli* (ATCC25922), *Klebsiella pneumoniae*, (ATCC700603) *Vibrio cholera*. (ATCC 39315) were procured from Ethiopian Health and Nutrition Research Institute (EHNRI) Clinical Bacteriology laboratory, Addis Ababa, Ethiopia.

Cultures of Bacteria

All the selected bacterial strains were cultured on nutrient agar plates and incubated for 24 hours at 37° C. The cultured bacterial strains were inoculated into Mueller-Hinton Broth and incubated at 37° C for 24 h before use.

Standardization of inoculate

 1×10^8 CFU/ml of the selected bacterial strains were used to test antibacterial activity of plant

extracts by using McFarland turbidity standard (Thirumurugan, 2010).

Antibacterial Activity (Agar Well-diffusion Assay)

The antibacterial activities of the crude leaf extracts of A. etbaica were determined by agar well diffusion method (Bauer *et al.*, 1966). 50μ l inoculums of tested bacterial strains were inoculated on the surface of solidified Mueller-Hinton Agar (MHA), by using a sterile cotton swab. The agar wells of 6-mm diameter were made using a sterile cork borer. About 100μ l of 10% solution (1 mg/ml) of each organic leaf extracts of *A. etbaica* in Dimethyl sulphoxide (DMSO) was dispensed in separate agar well with the help of a micropipette. The agar dishes were pre-incubated at 37°C for 2 hours to allow the uniform diffusion of organic leaf extracts of A. etbaica into the agar. After pre-incubation, the plates were incubated at 37°C for 24 hours. The dissolution of the organic leaf extracts of A. etbaica was facilitated with the addition of 10% (v/v) DMSO which not affected the growth of microorganisms (as shown by our control experiments).



Figure 3: Zone of inhibition (mm) of acetone leaf extract of *Acacia etbaica* (100, 250, 500 μ g/mL) and standard antibiotics Ciproflaxin (20 μ g/mL) against three gram negative and three gram positive bacteria by the Agar well-diffusion method

Besides Ciprofloxacin (20 μ g) was used a positive control for selected bacterial strains. The antibacterial activity was determined by measuring the inhibition zone diameter in millimeters in millimetre-scale after incubation (Lalitha, 2015).

Minimum inhibition concentration of effective plant extracts (MIC)

The acetone and ethanol leaf extract of *A. ethaica* were dissolved in 10% DMSO. The initial test concentrations were serially diluted in a 96 well plate and inoculated with 5 μ l of 10⁸ CFU ml⁻¹ bacterial suspensions which showed highly susceptible against the tested plant extract. The 96 well plates were incubated for 24 hours at 37 °C for bacterial growth. The culture intensity of each well was read at 600 nm and compared with the untreated controls. The MIC of the selected extract was determined as the lowest concentration of the extract inhibiting the visual growth of the test cultures (Wikler, 2009).

RESULTS AND DISCUSSION

Preliminary phytochemical screening

As shown in Table 1, the phytochemical analysis of

organic leaf extracts of *A. etbaica,* revealed that the acetone leaf extracts of *A. etbaica,* confirmed the presence of a maximum number of phytoconstituents like flavonoids, tannins, phenolic compounds terpenoids, and alkaloids were absent. The ethanolic leaf extracts of *A. etbaica* showed the presence of a significant number of phytoconstituents like flavonoids, tannins, phenolic compounds terpenoids and the absence of both alkaloids and coumarins. The current results were in agreement with previous reports of (Saleh *et al.*, 2014).

In vitro antibacterial activity of crude leaf extract of *A.etbaica* against *Enterobacteriaceae* family

The *in vitro* antibacterial activity of organic leaf extract of *A. etbaica* at various concentrations (100, 250, and 500μ g/ml) and standard antibiotic ciprofloxacin (20 μ g/ml) were shown in the (Tables 2, 3, 4, 5 and 6). The *A.etbaica* organic leaf extract showed the zone of inhibition (*in vitro* antibacterial activity) ranging from 10 mm to 18 mm diameter (diameters which included the 6 mm of agar wells) with the significant results were shown by both acetone and ethanolic leaf extract of *A. etbaica* against both selected bacterial strains of



Figure 4: Zone of inhibition (mm) of ethanol leaf extract of *Acacia ethaica* (100, 250, 500 μ g/mL) and standard antibiotics Ciproflaxin (20 μ g/mL) against three gram negative and three gram positive bacteria by the Agar well-diffusion method

Enterobacteriaceae family.

The antibacterial activity of petroleum ether leaf extract of *A. etbaica*, against the tested bacteria as shown in the Table 2 and Figure 1, exhibited insignificant zone of inhibition produced against the grampositive *E. faecalis* (10 mm) at 500 μ g/ ml and also showed same results with the gram-negative bacteria *V. cholerae* (9-13 mm) and *E. coli* (10-13 mm) at a concentration range between 100 μ g/ ml -500 μ g/ mL.

The antibacterial activity of chloroform leaf extract of *A. etbaica* as shown in Table 3 and Figure 2 exhibited prominent zone of inhibition produced against, tested gram-positive *B.subtilis* (11-12 mm), *S.aureus*(10-12 mm), and *E. faecalis* (10-12 mm) and the gram-negative bacteria *K. pneumoniae* (11-14mm), *V. cholera* (18- 20 mm) and *E. coli* (16- 24 mm) at a concentration range between 100 μ g/ ml -500 μ g/ mL.

The antibacterial activity of acetone leaf extract *A. etbaica*, as shown in Table 4 and Figure 3, exhib-

ited a significant broad spectrum of the zone of inhibition against the tested gram-negative bacteria *E. coli* (10- 17 mm), *V. cholera* (10-16 mm) and *K. pneumonia* (10- 17 mm) and the gram-positive bacteria *B.subtilis* (12- 16mm), *S.aureus* (12- 15mm) and *E. faecali* (11- 15 mm) at a concentration range between 100-500 μ g/ mL. Therefore acetone leaf extract *A. etbaica* showed an effective broad spectrum of *in vitro* antibacterial activity against tested both gram-positive and gram-negative bacteria.

The antibacterial activity of ethanolic leaf extracts *A. etbaica* as shown in Table 5 and Figure 4 had shown a broad spectrum of the highest zone of inhibition against gram-negative bacteria *E. coli* (11-18 mm), *V. cholera* (11-13 mm), *K. pneumoniae* (10-14 mm) and gram-positive bacteria *B.subtilis* (10-13mm), *S.aureus* (10-14mm) and *E. faecali* (10-14 mm) at 500 μ g/ ml at a concentration range between 250 μ g/ ml -500 μ g/ ml.

Minimal inhibitory concentration (MIC)

The Minimum Inhibitory Concentration (MIC) of the



Figure 5: Minimum Inhibitory Concentration (MIC) of acetone and ethanolic leaf extract of Acacia etbaica against Enterobactriace in μ g/ml

effective acetone and ethanolic leaf extract of A. etbaica (AE-AC and AE-ET) at various concentrations were assessed by 96 well plate broth microdilution methods with a concentration range from 1.25 μ g /ml to 100 μ g /ml. Only the tested bacteria, which were showed highly susceptible to the selected plant extracts was taken to determining the MIC. The maximal zones of inhibition and MIC values for tested bacterial strains, which were sensitive to the leaf extract of A. etbaica, were in the range of 10-18 mm and lowest MIC values of 0.532 μ g/ml against tested gram +ve bacteria. As showed in Table 5, the acetone leaf extract of A. etbaica was highly susceptible against tested *B.subtilis*, *S.aureus*, E. feacalis, V. cholea, K. pneumonia, and E.coli with the significant MIC value 0.532, 0.625, 2.5, 2.06, 4.12 and 15.60 μ g/ml respectively.

As showed in the Figure 5, the ethanolic leaf extract of *A. etbaica* was highly susceptible to *B.subtilis, S.aureus, E. feacalis, E.coli*, *V. cholea,* and *K. pneumonia,* with the significant MIC values of 1.24, 1.5, 4.12, 3.008, 14.86 and 24.87. Among the tested organic, leaf extract of *A. etbaica* both acetone leaf and ethanolic leaf extract of *A.etbaica* shown strong broad-spectrum antibacterial activity against tested Enterobactriace family in the range of 11- 18 mm with less MIC values of between 0.532 to 24.87 μg /ml.

CONCLUSIONS

Both the acetone and ethanolic leaf extract of Acacia etbaica exhibited a broad spectrum of promising in vitro antibacterial activities against tested MDR Enterobacteriaceae family of both selected bacterial strains among the other tested extracts. The qualitative phytochemical analysis of the acetone and ethanolic leaf extract of A ethaica shown the presence of the maximum number of phytoconstituents like flavonoids, tannins, terpenoids, and polyphenolic components. The high phenolic and flavonoid content present in both acetone and ethanolic leaf extract of A.etbaica accompanied by producing a significant zone of inhibition against all tested MDR Enterobacteriaceae family. The current studies demonstrated that the broad-spectrum antibacterial activity of acetone and ethanolic leaf extracts of Acacia etbaica may be helpful for the isolation of novel potent antibacterial agents against both hospital-acquired and communicable infectious bacterial pathogens without any side effects.

ACKNOWLEDGEMENT

The authors sincerely thank to the Department of Chemistry, Aksum University, and Aksum, Ethiopia for providing laboratory facilities.

Funding Support

This project work was partially funded by Ministry of Education, Ethiopia.

Conflict of Interest

The authors have confirm no conflict of interest.

REFERENCES

- Bauer, A. W., Kirby, W. M. M., Sherris, J. C., Turck, M. 1966. Antibiotic Susceptibility Testing by a Standardized Single Disk Method. *American Journal of Clinical Pathology*, 45(4_ts):493–496.
- Brenner, D. J., Farmer, J. F., Enterobacteriaceae 2005. Bergey's manual of systematic bacteriology. pages 587–606, New York. Springer.
- Das, K., Tiwari, Rks, Shrivastava, D. K. 2010. Techniques for evaluation of medicinal plant products as antimicrobial agent: Current methods and future trends. *Journal of Medicinal Plants Research*, 4(2):104–111.
- Fair, R. J., Tor, Y. 2014. Antibiotics and Bacterial Resistance in the 21st Century. *Perspectives in Medicinal Chemistry*, 6:PMC.S14459–PMC.S14459.
- Harborne, A. 1998. Phytochemical methods a guide to modern techniques of plant analysis. *springer science & business media*, 3rd:15–18.
- Jima, T. T., Megersa, M. 2018. Ethnobotanical Study of Medicinal Plants Used to Treat Human Diseases in Berbere District, Bale Zone of Oromia Regional State. pages 1–16.
- Kassem, A. S., Naser, A. A., Ahmed, A. M., Muna, T., Tariq, M. R. 2014. Anatomical and phytochemical studies of the leaves of Acacia etbaica subspecies etbaica. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 5(4):1–10.
- Lalitha, M. K. 2015. Christian Medical College. *Manual of Antimicrobial Susceptibility Testing*, pages 1–47.
- Melese, B., Paulos, W., Astawesegn, F. H., Gelgelu, T. B. 2019. Prevalence of diarrheal diseases and associated factors among under-five children in Dale District, Sidama zone, Southern Ethiopia: a cross-sectional study. *BMC Public Health*, 19(1):1235–1235.
- Munita, J. M., Arias, C. A. 2016. Mechanisms of Antibiotic Resistance. *Microbiology Spectrum*, 4(2):1–37.

Nikaido, H. 2009. Multidrug Resistance in Bacteria. *Annual Review of Biochemistry*, 78(1):119–146.

- Reta, A., Kifilie, A. B., Mengist, A. 2019. Bacterial Infections and Their Antibiotic Resistance Pattern in Ethiopia: A Systematic Review. *Advances in Preventive Medicine*, 2019:1–10.
- Ross, J. H. 1979. A Conspectus of the African Acacia Species. Memoirs of the Botanical Survey of South Africa No. 44. Botanical Research Institute, Department of Agricultural Technical Services. Republic of South Africa.
- Saleh, A., Algifri, K., Mohammed, A. N., Taleb, A., Ramzi, M. T. M. 2014. Anatomical and Phytochemical Studies of the Leaves of Acacia etbaica subspecies Etbaica. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 5(4):802– 810.
- Teklay, A., Abera, B., Giday, M. 2013. An ethnobotanical study of medicinal plants used in Kilte Awulaelo District, Tigray Region of Ethiopia. *Journal of Ethnobiology and Ethnomedicine*, 9(1):65–65.
- Thirumurugan, K. 2010. Antimicrobial activity and phytochemical analysis of selected Indian folk medicinal plants. *Steroids*, 1(10):430–434.
- Wikler, M. 2009. Clinical and Laboratory Standards Institute Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard. volume 35, pages 9–110, Wayne, USA.