



## Face and content validation of Phacoemulsification Techniques Related to Surgically Induced Astigmatism questionnaire

Md Muziman Syah Md Mustafa<sup>\*1</sup>, Nazaryna Marzuki<sup>2</sup>, Khairidzan Mohd Kamal<sup>3</sup>,  
Noorhazayti Ab. Halim<sup>4</sup>

<sup>1</sup>Department of Optometry and Visual Science, Kulliyah of Allied Health Sciences, International Islamic University Malaysia, Kuantan, Pahang, Malaysia

<sup>2</sup>Department of Ophthalmology, Sarawak General Hospital, Ministry of Health Malaysia, Kuching, Sarawak, Malaysia

<sup>3</sup>Department of Ophthalmology, Kulliyah of Medicine, International Islamic University Malaysia, Kuantan, Pahang, Malaysia

<sup>4</sup>Department of Paediatric Dentistry and Dental Public Health, Kulliyah of Dentistry, International Islamic University Malaysia, Kuantan, Pahang, Malaysia

### Article History:

Received on: 15.03.2019

Revised on: 20.06.2019

Accepted on: 25.06.2019

### Keywords:

Face validity,  
Content validity,  
Phacoemulsification,  
Questionnaire,  
Surgically induced  
astigmatism

### ABSTRACT

Phacoemulsification surgical techniques are associated with surgically induced astigmatism (SIA) in post-cataract surgery. However, no established tool is currently available to gather information on the surgeon's phacoemulsification techniques. This study aimed to validate a newly developed 'Phacoemulsification Techniques Related to Surgically Induced Astigmatism' (PTechSIA) questionnaire. PTechSIA questionnaire was designed to collect information on surgical techniques used by surgeons during phacoemulsification. Questionnaire items were generated based on peer-reviewed literature of related domains to phacoemulsification surgical techniques. The self-administered tool consisted of fifteen close-ended questions. Ten subject-matter experts were recruited for face validation using qualitative approach and content validation using Lawshe's method. Content validation ratio (CVR) was calculated for each item; items with CVR value of less than 1.00 were eliminated, and the final content validity index (CVI) of the questionnaire was obtained. Out of fifteen items, six items with CVR value of less than 1.00 were eliminated with initial CVI of 0.61. After the modification process, the number of final items was nine, with an overall final CVI score of 1.00. Hence, the findings indicate that the PTechSIA questionnaire can be utilized for future assessments of surgical techniques and their association with SIA.



### \*Corresponding Author

Name: Md Muziman Syah Md Mustafa

Phone:

Email: syah@iiium.edu.my

ISSN: 0975-7538

DOI: <https://doi.org/10.26452/ijrps.v10i4.1531>

Production and Hosted by

IJRPS | [www.ijrps.com](http://www.ijrps.com)

© 2019 | All rights reserved.

### INTRODUCTION

A qualified measuring tool is defined by the reliability and validity of the measures. A researcher must evaluate the validity of an assessment tool before its implementation (Mohamed *et al.*, 2017) in order to minimise measurement errors (Gilbert and Prion, 2016). Content validation of a measuring tool is referred to as the extent of the tool which covers all the domains being studied. The contents should be extracted from literature, and suggestions were made and evaluated by at least three subject-matter

experts (SMEs) (Lynn, 1986).

Surgically induced astigmatism (SIA) in post phacoemulsification cataract surgery is determined by the surgical techniques. Several surgical techniques associated with SIA had been discussed in previous studies (Kawahara *et al.*, 2013; Kwon *et al.*, 2014; Chang *et al.*, 2015; Nikose *et al.*, 2018). Other factors such as the location and size of the main incision are also frequently reported as significant factors that are related to the SIA, although not regularly documented in surgical notes. There are also other surgical techniques which are often neglected; these include hand dominance, microkeratome design, as well as design and type of incisions which are possibly associated with SIA.

Although there were several researches conducted about the surgical techniques related to SIA, no specific tool was developed in documenting all the surgical techniques used in phacoemulsification. Hence, it is necessary to have a reliable and validated questionnaire for this purpose.

## MATERIALS AND METHODS

This study obtained ethical approval from the International Islamic University Malaysia (IIUM) Research Ethical Committee (Reference Number: IREC 2018-65).

### Tool Development

The tool was initially developed based on peer-reviewed literature. The first items included several domains of phacoemulsification techniques which were related to SIA. The domains were 1) the main incision, 2) the side port incision, 3) the position of surgeon, 4) the phacoemulsification technique, and 5) the experience of the surgeon (Kawahara *et al.*, 2013; Yoon *et al.*, 2013; Theodoulidou *et al.*, 2015).

### Face and Content Validation

Ten subject-matter experts (SMEs) who are qualified Ophthalmologists from the Ministry of Health, Malaysia ( $n = 10$ ) participated in this study. Five SMEs were involved in face validation, whereby the other five were involved in content validation. Each SME was given a copy of the questionnaire with a research information sheet and consent form. The purposes, benefits, risks and procedures of this study were briefly explained to all SMEs.

### Face Validity

Face validation is a process in which the target group rates a test item, whether it is relevant or does not fit its purpose (Devon *et al.*, 2007). A qualitative approach was used to conduct a face validity assessment (Sangoseni *et al.*, 2013; Abootalebi

*et al.*, 2016). The five SMEs assessed each item in the questionnaire thoroughly by looking at the style and format consistency, language clarity and readability (Devon *et al.*, 2007), sentence syntax and practicality, and suitability of terminologies used. The SMEs also provided feedback on the difficulties and ambiguities encountered while filling out the questionnaire. Self-administered responses were recorded directly in the "comment/suggestion" section of the questionnaire.

### Content Validity Ratio and Content Validity Index

Content validity ratio (CVR) was evaluated using Lawshe's method. The CVR is an established statistical method to determine the content validity of individual items by retaining or rejecting the items (Wilson *et al.*, 2012) using the following equation:

$$CVR = (n_e - N/2) \div (N/2)$$

$n_e$  = a number of SMEs rating a measurement item as "essential".

$N$  = the total number of SMEs who are involved in the content validity process.

The CVR value ranges from -1.00 to 1.00. Scores of 0.00 to 0.99 indicates that more than half but less than all SMEs rated an item as "essential" while rating 1.00 suggests that all of SMEs rated a thing as "essential". A negative CVR value indicates that more than half of SMEs did not rate the items as "essential". The decision on item acceptance was based on Lawshe's table (Lawshe, 1975) containing threshold values, in which only the items with CVR above the threshold value would be accepted. Lawshe's threshold value is different depending on the number of SMEs involved. For this current study, which recruited five SMEs, the threshold CVR value was 0.99. In other words, it requires all SMEs to agree that an item is considered "essential" for it to be accepted in the final tool.

Content validity of the entire questionnaire was evaluated using the content validity index (CVI). The CVI is the mean of CVR values of all individual items. CVI value of more than 0.80 shows good content validity of the overall questionnaire (Davis, 1992).

## RESULTS AND DISCUSSION

SIA has been emphasized in previous studies as it leads to better visual performance and satisfaction in post-cataract surgery (Gundersen and Potvin, 2016; Srivannaboon and Chirapapaisan, 2017; Clark, 2018). The surgical technique is an established factor that influences SIA (Kawahara *et al.*, 2013; Kwon *et al.*, 2014; Chang *et al.*, 2015; Nikose *et al.*, 2018). However, incomplete infor-

**Table 1: Content validation assessment**

Domains	Item Number	Items	CVR	Decision	
Main incision	1	Which one is your dominant hand?	1.00	Accept	
	3	What is your main incision architecture?	0.50	Reject	
	4	What is the type of the main incision that you frequently perform?	1.00	Accept	
	6	What is the size of your main incision?	1.00	Accept	
	7	Do you measure the size of the main incision?	0.50	Reject	
	8	Where is the location of your main incision?	1.00	Accept	
	9	Do you have any specific microkeratome design in performing the main incision?	1.00	Accept	
	Side port incision	10	How many side port incision(s) do you perform?	1.00	Accept
		11	What is the size of your side port incision(s)?	0.50	Reject
12		Where is the location of your side port incision(s) in relative to the main incision?	1.00	Accept	
13		Do you mark the side port incision(s) location before surgery?	0.00	Reject	
14		Do you use any specific microkeratome design in performing the side port incision(s)?	0.50	Reject	
Surgeon's position	2	Where do you position yourself during the surgery?	0.50	Reject	
Phacoemulsification technique	5	What is the surgical technique that you apply?	1.00	Accept	
Surgeon's experience	15	How many phacoemulsification cataract surgery cases that you have performed?	0.50	Reject	

CVR - Content validation ratio

mation on surgical techniques in operative notes could hinder retrospective studies of SIA. It is suggested that surgical operation notes must have detailed surgery procedures, all operative findings and postoperative instructions to enable continuity of care by other ophthalmic surgeons. Therefore, the patient's operation note should be completed and legible for future readers (Payne *et al.*, 2011). As to date, there are no standardized templates for oper-

ation notes related to SIA that have been published. This current PTechSIA questionnaire was aimed to be an initial milestone towards comprehensive and legitimate documentation as a solution to the missing essential surgical information. To initiate the process, fifteen items were generated which consisted of seven items for the main incision domain, five items for the side port incision domain, and one item for the position of surgeon, phacoemulsifica-

tion technique and surgeon's experience domains, respectively. Subsequently, the first fifteen items underwent face and content validation processes by SMEs before being used in clinical or research settings.

The purpose of validation exercise is to identify any unrelated items that do not fit the studied domain and to amend the items for improvement (Lynn, 1986). In this current study, face and content validation were performed to identify and filter the non-fit items. Several conflicting reports were found in previous studies regarding the definition and assessment method of face validity (Cook and Beckman, 2006; Bölenius et al., 2012). There was no consensus on the status, role and importance in executing face validity assessment. However, we believed that face validity is an important test to confirm the tool appropriateness, comprehension and measures the questionnaire objectives (Bölenius et al., 2012). We conducted a qualitative approach in the face validity process in accordance with several previous works (Sangoseni et al., 2013; Abootalebi et al., 2016). We found that the qualitative approach enables SMEs to provide subjective analysis for each item and give their expert opinion for the improvement of the questionnaire.

Modifications of the questionnaire were performed according to the suggestions given by SMEs. The modified version of the questionnaire underwent further content validation process. There are several approaches in content validation, as explained in previous literature (Lawshe, 1975; Polit et al., 2007; Gilbert and Prion, 2016). Lawshe's method was chosen in this current study due to its wide acceptance and simplicity (Ayre and Scally, 2014). Due to the absence of reports on ways to calculate original critical values, later studies have attempted to revise the critical value of Lawshe's CVR (Wilson et al., 2012; Ayre and Scally, 2014), which revealed anomalous values in the original Lawshe's CVR critical value. However, the anomaly does not affect the stringency of the value, and it is a safe method to use in deciding the inclusion of items (Wilson et al., 2012). In this current study, the revised critical values of the five SMEs were similar between the original and revised versions. Therefore, it can be confirmed that the accepted items were appropriately content validated. The assessment summary is displayed in Table 1.

Overall CVI score of the questionnaire with all fifteen items was 0.61. Then, the following six items (items number 2, 7, 11, 13, 14, 15) with CVRs of less than 0.99 were excluded from the questionnaire. After modification of the questionnaire by

retaining nine items, the overall final CVI score of 1.00 was obtained. Previous literature outlined that CVI value of more than 0.80 indicates good agreement (Davis, 1992) and 1.00 represents a perfect agreement between SMEs (Lynn, 1986). This proves that content validity for the entire questionnaire is acceptable where all the SMEs achieved consensus in their assessments.

## CONCLUSION

Proper validation processes are crucial in the development of a questionnaire as a research and clinical tool. In this research, the PTechSIA questionnaire has passed through a proper face and content validity processes. To the best of our knowledge, there is no published tool related to SIA currently available. Therefore, the PTechSIA questionnaire can be utilized in the near future to assess surgical techniques and their association with SIA. Moreover, it could benefit clinician and researcher who intend to perform a further investigation of the association of surgical techniques and SIA.

## ACKNOWLEDGEMENT

Special thanks to IIUM research funding for financially supporting this research (RIGS 16-129-0293).

## REFERENCES

- Abootalebi, M., Kargar, M., Jahanbin, I., Sharifi, A. A., Sharafi, Z. 2016. Knowledge and attitude about andropause among general physicians. *International Journal Community Based Nursing Midwifery*, 4(1):27-35.
- Ayre, C., Scally, A. J. 2014. Critical values for Lawshe's content validity ratio. *Measurement and Evaluation in Counseling and Development*. 47:79-86.
- Bölenius, K., Brulin, C., Grankvist, K., Lindkvist, M., Söderberg, J. 2012. A content validated questionnaire for assessment of self-reported venous blood sampling practices. *BMC Research Notes*, 5(1):39.
- Chang, S. W., Su, T. Y., Chen, Y. L. 2015. Influence of ocular features and incision width on surgically induced astigmatism after cataract surgery. *Journal of Refractive Surgery*, 31(2):82-88.
- Clark, K. D. 2018. Toric intraocular lens outcomes with a new protocol for IOL selection and implantation. *Journal Francais D'Ophthalmologie*, 41(2):145-151.
- Cook, D. A., Beckman, T. J. 2006. Current concepts in validity and reliability for psychometric instruments: theory and application. *The American Journal of Medicine*, 119(2):166-173. e16.

- Davis, L. L. 1992. Instrument review: Getting the most from a panel of experts. *Applied Nursing Research*, 5(4):194–197.
- Devon, H. A., Block, M. E., Moyle-Wright, P., Ernst, D. M., Hayden, S. J., Lazzara, D. J., Kostas-Polston, E. 2007. A psychometric toolbox for testing validity and reliability. *Journal of Nursing Scholarship*, 39(2):155–164.
- Gilbert, G. E., Prion, S. 2016. Making sense of methods and measurement: Lawshe's content validity index. *Clinical Simulation in Nursing*, 12:530–531.
- Gundersen, K. G., Potvin, R. 2016. Clinical outcomes with toric intraocular lenses planned using an optical low coherence reflectometry ocular biometer with a new toric calculator. *Clinical Ophthalmology*, 10:2141–2147.
- Kawahara, A., Kurosaka, D., Yoshida, A. 2013. Comparison of surgically induced astigmatism between one-handed and two-handed cataract surgery techniques. *Clinical Ophthalmology*, 7:1967–1972.
- Kwon, H. J., Nam, S. M., Stulting, R. D., Lim, C. Y., Seo, K. Y. 2014. Comparison of surgically induced astigmatism following iris-claw PIOL insertion with scleral, limbal, or corneal incisions. *Journal of Refractive Surgery*, 30(5):330–335.
- Lawshe, C. 1975. A quantitative approach to content validity. *Personnel Psychology*, 28(4):563–575.
- Lynn, M. 1986. Determination and quantification of content validity. *Nursing Research*, 35(6):382–385.
- Mohamed, S. T. S., Reddy, K., Chakka, G., Kumar, B. M., Reddy, S. V. 2017. Assessment of medication usage awareness among rural population living in resource limited settings. *International Journal of Research in Pharmaceutical Sciences*, 8(4):621–628.
- Nikose, A. S., Saha, D., Laddha, P. M., Patil, M. 2018. Surgically induced astigmatism after phacoemulsification by temporal clear corneal and superior clear corneal approach: a comparison. *Clinical Ophthalmology*, 12:65–70.
- Payne, K., Jones, K., Dickenson, A. 2011. Improving the standard of operative notes within an oral and maxillofacial surgery department, using an operative note proforma. *Journal of Maxillofacial and Oral Surgery*, 10(3):203–208.
- Polit, D., Beck, C., Owen, S. 2007. Is the CVI an acceptable indicator of content validity? Appraisal and recommendations. *Research in Nursing and Health*, 30(4):459–467.
- Sangoseni, O., Hellman, M., Hill, C. 2013. Development and validation of a questionnaire to assess the effect of online learning on behaviors, attitudes, and clinical practices of physical therapists in the United States regarding evidenced-based clinical practice. *The Internet Journal of Allied Health Sciences and Practice*, 11:231–235.
- Srivannaboon, S., Chirapapaisan, C. 2017. Consistency analysis of surgically induced astigmatism. *Journal of Cataract and Refractive Surgery*, 43(8):1117–1118.
- Theodoulidou, S., Asproudis, I., Kalogeropoulos, C., Athanasiadis, A., Aspiotis, M. 2015. The role of sideport incision in astigmatism change after cataract surgery. *Clinical Ophthalmology*, 9:1421–1428.
- Wilson, F. R., Pan, W., Schumsky, D. A. 2012. Recalculation of the critical values for Lawshe's content validity ratio. *Measurement and Evaluation in Counseling and Development*. 45:197–210.
- Yoon, J. H., Kim, K. H., Lee, J. Y., Nam, D. H. 2013. Surgically induced astigmatism after 3.0 mm temporal and nasal clear corneal incisions in bilateral cataract surgery. *Indian Journal of Ophthalmology*, 61(11):645–648.