



Combined Treatment with Oral Hypoglycemic Agents and Insulin in Longer Run May Lead to Cognitive Derangement Secondary to Hypoglycemia

Aman Gupta^{*1}, Abhishek Kumar Singh¹, Ramesh Deka C²

¹Amity Institute of Neuropsychology and Neurosciences, Amity University Uttar Pradesh, Sector-125, Noida-201313, India

²Department of Medical Education & Research, Amity University Uttar Pradesh India, Previously Director, All India Institute of Medical Sciences, Ansari Nagar, New Delhi-110029, India

Article History:

Received on: 24.09.2019

Revised on: 15.12.2019

Accepted on: 20.12.2019

Keywords:

Diabetes,
diabetes complications,
OHA,
Insulin therapy,
hypoglycemia

ABSTRACT

To determine the risk associated with combined therapy of oral hypoglycemic agents and insulin in the development of cognitive decline secondary to hypoglycemia. The study comprised of two arms, in arm 1 (n=30) diabetic patients on OHA and insulin combined therapy were observed for cognitive derangement based on Mini Cog score (Pearson's correlation r was calculated between the two) and the patients were also ruled out for hypoglycemia episodes. Similarly, in arm 2(n=30), only OHA therapy patients were evaluated on the same parameters using Pearson's correlation. Mean Mini Cog scores were found to be 1.79 in patients on combinational therapy of OHA and insulin. In the case of patients only on OHA for diabetes control mean Mini Cog, the score was 2.0. Pearson's coefficient r 1 between OHA plus insulin duration of therapy in years with Mini Cog score was -0.5831 at a statistically significant p value of 0.0007. While on the other hand, patients only on OHA therapy (in years) were not significantly correlated with Mini Cog scores.(r 2= - 0.0149 ; p value =0.93771).In the arm 1 (OHA + Insulin(n=30)), the average number of hypoglycemia episodes were found to be 2.633 as compared to the OHA only group in which mean hypoglycemia episodes were 0.866. There was a significant difference in the hypoglycemia episodes, as confirmed by the student t-test (4.06368) at a p value = 0.000147. Based on the study findings, combined Treatment with Oral Hypoglycemic agents and insulin in longer run may lead to cognitive derangement secondary to hypoglycemia.*Significance level <0.01, Negative correlations are indicative of lower the values of battery scores higher would be cognitive derangement.



*Corresponding Author

Name: Aman Gupta

Phone: 919953270158

Email: gupta.draman@gmail.com

ISSN: 0975-7538

DOI: <https://doi.org/10.26452/ijrps.v11i1.1852>

Production and Hosted by

IJRPS | www.ijrps.com

© 2020 | All rights reserved.

INTRODUCTION

Diabetes is one of the key disease burdens of the globe. Multiple studies have been conducted to find an association between diabetes and cognitive decline and dementia. Oral hypoglycemic agents are the backbone for managing type 2 Diabetes in an early phase (ICMR, 2018). Sulphonylureas are associated with hypoglycemia episodes (Klein-Schwartz et al., 2016). In a metanalysis (Andersen and Christensen, 2016) did find adding more than one therapy to OHA may lead to hypoglycemia. In another study, it was found that adding insulin to met-

formin can have a higher risk of developing hypoglycemia (Roumie *et al.*, 2016). Research suggests that the risk of hypoglycemia increases with age, duration of Diabetes and combinational therapy of OHA and Insulin (Amiel *et al.*, 2008).

In a retrospective study conducted in Japan, multiple factors like age, cognitive decline, combinational therapy with of Insulin and sulphonylureas have been cited as risk factors for severe hypoglycemia (Ikeda *et al.*, 2018).

In a study conducted by (Lin *et al.*, 2018), about 66% of the patients on combination therapy of OHA and Insulin suffered with moderate hypoglycemia (Chin *et al.*, 2016) investigated the association between hypoglycemia episodes and cognitive decline and did find that linear correlation between the two. In another study, it was found that in the case of severe hypoglycemia episodes, the risk of development of cognitive decline was much higher (Ryan *et al.*, 2016). Another aspect to be kept in consideration with respect to the pathophysiology of cognitive decline is advanced glycaemic end products (AGEs), which are produced secondary to uncontrolled long-duration diabetes (Gorska-Ciebiada *et al.*, 2015). Brain insulin resistance has also been found correlated with cognitive decline in Diabetes (Arnold *et al.*, 2018).

Cerebral microinfarcts also add to the pathophysiology of dementia in Diabetes type 2 (Biessels and Despa, 2018).

(Rasch and Born, 2013) explained the importance of sleep in memory consolidation. Others indicated REM sleep is more important for memory consolidation.

(Gupta *et al.*, 2018) presented a correlation between different pathophysiologicals involved in the cognitive derangement in Diabetes (Gupta, 2018) and role of sleep in memory consolidation and mechanism of cognitive decline in case of poor sleep or decreased sleep (Refer Figure 1).

In the current study, we hypothesized that combinational OHA and Insulin therapy has a higher risk of developing cognitive derangement as compared to OHA only group. We further hypothesized that higher cognitive decline in the OHA plus Insulin therapy group can be secondary to hypoglycemia as the combinational therapy had a higher incidence of hypoglycemia episodes.

MATERIALS AND METHODS

Patients were enrolled from either outpatient department or hospitalized patients from Sarvodaya Hospital and Research Center, Faridabad, Delhi

NCR, India. Ethical clearance was sought from the University Ethics Committee of Amity University Uttar Pradesh, Noida, All the subjects have provided written informed consent to participate in the study.

The study comprised of two arms (refer Figure 6), in arm 1 (n=30) diabetic patients on OHA and insulin combined therapy were observed for cognitive derangement based on Mini Cog score (Pearson's correlation r was calculated between the two) and the patients were also ruled out for hypoglycemia episodes. Similarly, in arm 2 (n=30), only OHA therapy patients were evaluated on the same parameters using Pearson's correlation. Demographics details can be referred to as per the Table 1. The current study was a single point prevalence study.

Inclusion and Exclusion criteria

Inclusion criteria

1. 35 -80 years of age
2. Type 2 DM patients as per "American Diabetic Association "
3. Greater than equal to 5 years of Diabetic History
4. Greater than 7 HBA1C
5. Willing to participate in the study and provide informed consent

Exclusion criteria

1. Age less than 35 years
2. Pregnant women and Children
3. Substance abuse
4. Significant Neurological, Psychiatric disease

Cognitive batteries

We used Mini Cog battery for cognitive assessment post getting written permission from

"The Alzheimer's Association" (<http://www.alz.org>).

Statistical analysis

We used IBM SPSS trial version for the statistical analysis. Mini Cog depicted the cognitive decline of the patients. To establish an association between Mini Cog and duration of OHA plus Insulin therapy in years, Pearson's correlation coefficient (r) was used. Similarly, Pearson's correlation coefficient (r) was

Table 1: Different variables along with the respective range

Variables	Range
Age (Years)	35-75
Weight (kg)	45-92
Duration of diabetes (Years)	6-17
No. of hypoglycemic episodes	0-9
HBA1C values	7-17

Table 2: OHA+Insulin therapy versus OHA treatment groups versus corresponding Pearson’s coefficient

Treatment	Group- Therapy duration in years	Mean Mini Cog Score	Pearson’s Correlation	p value (<0.05)
OHA + Insulin(n=30)		1.79	-0.5831	0.0007*
OHA only (n=30)		2	-0.0149	0.93771**
		*Significant	**Not Significant	

Table 3: OHA+Insulin therapy versus OHA treatment groups versus corresponding Hypoglycemia episodes in both arms

Treatment	Group- Therapy duration in years	Mean number Hypoglycemia Episodes	t-test value	p value(<0.05)
OHA + Insulin(n=30)		2.633333333	4.06368	0.000147* *Significant




Figure 1: Proposed hypothesis




Figure 2: Box and Whisker Plot -OHA +Insulin Therapy(years)versus Mini-Cog Score




Figure 3: Box and Whisker Plot -only OHA Therapy (years)versus Mini-Cog Score

also calculated for the arm on only OHA therapy to find an association with Mini Cog scores.

Both the groups were also assessed for hypoglycemia episodes and the t-test was used to compare the results of both the arms at a p value less than 0.05 and 95% confidence interval.

RESULTS AND DISCUSSION

OHA and Insulin combinational therapy was associated with higher Cognitive derangement as compare to OHA only arm

Mean Mini Cog scores were found to be 1.79 in patients on combinational therapy of OHA and

insulin. In the case of patients only on OHA for diabetes control, mean Mini- Cog score was 2.0. Pearson’s coefficient r_1 between OHA plus insulin duration of therapy in years with Mini Cog score was -0.5831 at a statistically significant p value of 0.0007 . While on the other hand, patients only on OHA therapy (in years) were not significantly correlated with Mini Cog scores. ($r_2 = -0.0149$; p value $=0.93771$). Refer to Table 2 for details.

Higher Cognitive derangement in OHA and Insulin combinational therapy may be associated with hypoglycemia episodes in this arm

In the arm 1 (OHA + Insulin(n=30)), the average number of hypoglycemia episodes were found to




Figure 4: Scatter Plot -OHA +Insulin Therapy (years)versus Mini-Cog Score




Figure 5: Scatter Plot -only OHA Therapy (years)versus Mini-Cog Score

be 2.633 as compared to the OHA only group in which mean hypoglycemia episodes were 0.866. There was a significant difference in the hypoglycemia episodes, as confirmed by the student t-test (4.06368) at a p value = 0.000147. Refer to Table 3 for details.

Multiple studies have been conducted to find an association between Diabetes and cognitive derangement, which have been bringing forward concepts pertaining to the insulin resistance and dementia. memory impairment (Schrijvers *et al.*, 2010). Another group did find the relation between Alzheimer’s disease and diabetes (Akter *et al.*, 2011). Researchers (Palta *et al.*, 2018) proposed its good to have diabetes in control to reduce the risk of cognitive decline. Animal models have also been successful to build up relationship between diabetes and dementia (Chomova *et al.*, 2017).

In the current study, we hypothesized that combinational OHA and Insulin therapy has a higher risk of developing cognitive derangement as compared to OHA only group. We further hypothesized that higher cognitive decline in the OHA plus Insulin therapy group can be secondary to hypoglycemia as the combinational therapy had a higher incidence of hypoglycemia episodes.

Current study results revealed that mean Mini Cog scores to be 1.79 in patients on combinational therapy of OHA and insulin and the same was 2.0 for the OHA only arm. Pearson’s coefficient r 1 between OHA plus insulin duration of therapy in years with Mini Cog score was -0.5831 at a statistically significant p value of 0.0007. While on the other hand, patients only on OHA therapy (in years)were not significantly correlated with Mini Cog scores.(r 2= - 0.0149 ; p value =0.93771).Figure 2 depicts Box and Whisker Plot -only OHA plus




Figure 6: Describe the study pattern and distribution between the two arms

Insulin Therapy (years) versus Mini-Cog Score and Figure 3 indicate Box and Whisker Plot -only OHA Therapy (years) versus Mini-Cog Score.

With reference to Figure 4 Scatter Plot -OHA plus Insulin Therapy (years) versus Mini-Cog Score and Figure 5 depicts Scatter Plot -only OHA Therapy (years) versus Mini-Cog Score.

We can mark the difference in the presentation in both the cases. These results are in line with earlier studies, which Research suggests that the risk of hypoglycemia increases with age, duration of Diabetes and combinational therapy of OHA and Insulin (Amiel *et al.*, 2008).

With respect to hypoglycemia episodes, in the arm 1 (OHA + Insulin (n=30)), the average number of hypoglycemia episodes were found to be 2.633 as compared to the OHA only group in which mean hypoglycemia episodes were 0.866. There was a significant difference in the hypoglycemia episodes, as confirmed by the student t-test (4.06368) at a p

value = 0.000147. These results are also aligned with previous studies like a retrospective study conducted in Japan, which indicated multiple factors like age, cognitive decline, combinational therapy with of Insulin and Sulphonylureas have been cited as risk factors for severe hypoglycemia (Ikeda *et al.*, 2018).

Different studies have provided evidence of cognitive decline secondary to hypoglycemia episodes. (Chin *et al.*, 2016) investigated the association between hypoglycemia episodes and cognitive decline and did find that linear correlation between the two. In another study, it was found that in the case of severe hypoglycemia episodes, the risk of development of cognitive decline was much higher (Ryan *et al.*, 2016).

CONCLUSIONS

The current study provides significant findings related to the optimal usage of drugs for the man-

agement of type 2 diabetes by virtue of its results. To our knowledge, no other similar study has been conducted in north India. Cognitive assessment was based on the Mini-Cog battery, which has already been validated.

We compared two arms of the study: OHA plus insulin therapy and only OHA treatment based on the correlation of individual arms with Mini Cog scores. We further evaluated, an association of these two categories of patients with a probability of hypoglycemia episodes and their significance levels. Based on the study findings, combined Treatment with Oral Hypoglycemic agents and insulin in the longer run may lead to cognitive derangement secondary to hypoglycemia.

We studied multiple aspects related to cognitive decline and diabetes. Uncontrolled Diabetes has been associated with cognitive derangement and so is hypoglycemia episodes. OHA and Insulin therapy, in combination, has a higher risk of hypoglycemia episodes, later, in turn, can increase the probability of cognitive decline.

Combinational therapy with OHA and insulin can be necessary to manage the poorly controlled diabetes. Hence, it is very important to keep track of the treatment strategy while managing uncontrolled diabetes. Further, large scale studies are required to further validate the concept. Randomized double-blind placebo-controlled clinical trials can provide further support to current evidence.

ACKNOWLEDGEMENTS

We thank "The Alzheimer's Association" (<http://www.alz.org>) for support with respect to usage of Mini COG used in the study.

REFERENCES

- Akter, K., Lanza, E. A., Martin, S. A., Myronyuk, N., Rua, M., Raffa, R. B. 2011. Diabetes mellitus and Alzheimer's disease: shared pathology and treatment? *British Journal of Clinical Pharmacology*, 71(3):365-376.
- Amiel, S. A., Dixon, T., Mann, R., Jameson, K. 2008. Hypoglycemia in Type 2 diabetes. *Diabetic Medicine*, 25(3):245-254.
- Andersen, S. E., Christensen, M. 2016. Hypoglycemia when adding sulphonylurea to metformin: a systematic review and network meta-analysis. *British Journal of Clinical Pharmacology*, 82(5):1291-1302.
- Arnold, S. E., Arvanitakis, Z., Macauley-Rambach, S. L., Koenig, A. M., Wang, H. Y., Ahima, R. S., Nathan, D. M. 2018. Brain insulin resistance in type 2 diabetes and Alzheimer disease: concepts and conundrums. *Nature Reviews Neurology*, 14(3):168-181.
- Biessels, G. J., Despa, F. 2018. Cognitive decline and dementia in diabetes mellitus: mechanisms and clinical implications. *Nature Reviews Endocrinology*, 14(10):591-604.
- Chin, S. O., Rhee, S. Y., Chon, S., Baik, S. H., Park, Y., Nam, M. S., Kim, Y. S. 2016. Hypoglycemia is associated with dementia in elderly patients with type 2 diabetes mellitus: An analysis based on the Korea National Diabetes Program Cohort. *Diabetes Research and Clinical Practice*, 122:54-61.
- Chomova, M., Balazova, M., Muchova, J. 2017. Diabetes-induced abnormalities of mitochondrial function in rat brain cortex: the effect of n-3 fatty acid diet. *Molecular and Cellular Biochemistry*, 435(1-2):109-131.
- Gorska-Ciebiada, M., Saryusz-Wolska, M., Borkowska, A., Ciebiada, M., Loba, J. 2015. C-Reactive Protein, Advanced Glycation End Products, and Their Receptor in Type 2 Diabetic, Elderly Patients with Mild Cognitive Impairment. *Frontiers in Aging Neuroscience*, 7.
- Gupta, A. 2018. Understanding common pathophysiologies associated with dementia and cognitive decline in sleep deficiency and diabetes. *Clin Psychiatry*, 4:12-12.
- Gupta, A., Singh, A., Deka, R. C., Gupta, R., Jha, R. 2018. To Investigate the Role of Glycosylated Hemoglobin (Hba1c) as a Biomarker for Prediction of Dementia and Cognitive Dysfunction in Type 2 Diabetic Patients. *Journal of Alzheimer's Disease & Parkinsonism*, (02):437-437.
- ICMR 2018. Guidelines for Management of Type 2 Diabetes. pages 23-26.
- Ikeda, Y., Kubo, T., Oda, E., Abe, M., Tokita, S. 2018. Incidence rate and patient characteristics of severe hypoglycemia in treated type 2 diabetes mellitus patients in Japan: Retrospective Diagnosis Procedure Combination database analysis. *Journal of Diabetes Investigation*, 9(4):925-936.
- Klein-Schwartz, W., Stassinis, G. L., Isbister, G. K. 2016. Treatment of sulphonylurea and insulin overdose. *British Journal of Clinical Pharmacology*, 81(3):496-504.
- Lin, S., Chen, M., Chen, W., Lin, K., Mu, P., Zhu, B., Zeng, L. 2018. A Randomized Trial of Insulin Glargine plus Oral Hypoglycemic Agents versus Continuous Subcutaneous Insulin Infusion to Treat Newly Diagnosed Type 2 Diabetes. *Journal of Diabetes Research*, pages 1-9.
- Palta, P., Carlson, M. C., Crum, R. M., Colantuoni, E.,

- Sharrett, A. R., Yasar, S., Golden, S. H. 2018. Diabetes and Cognitive Decline in Older Adults: The Ginkgo Evaluation of Memory Study. *The Journals of Gerontology: Series A*, 73(1):123-130.
- Rasch, B., Born, J. 2013. About Sleep's Role in Memory. *Physiological Reviews*, 93(2):681-766.
- Roumie, C. L., Min, J. Y., Greevy, R. A., Grijalva, C. G., Hung, A. M., Liu, X., Griffin, M. R. 2016. Risk of hypoglycemia following the intensification of metformin treatment with insulin versus sulfonylurea. *Canadian Medical Association Journal*, 188(6):104-112.
- Ryan, C. M., Klein, B. E. K., Lee, K. E., Cruickshanks, K. J., Klein, R. 2016. Associations between recent severe hypoglycemia, retinal vessel diameters, and cognition in adults with type 1 diabetes. *Journal of Diabetes and Its Complications*, 30(8):1513-1518.
- Schrijvers, E. M. C., Witteman, J. C. M., Sijbrands, E. J. G., Hofman, A., Koudstaal, P. J., Breteler, M. M. B. 2010. Insulin metabolism and the risk of Alzheimer disease: The Rotterdam Study. *Neurology*, 75(22):1982-1987.